

# Studies on the Effect of Different Combinations of Soybean Straw and Sugarcane Bagasse on Yield of Oyster Mushroom

P. S. Nagre, A. C. Jadhav, B. B. Chirame, C. T. Kumbhar and A. A. Bhagat  
All India Coordinated Research Project on Mushroom, College of Agriculture, Pune  
Mahatma Phule Krishi Vidyapeeth, Rahuri - 413 722 (India)  
(Received : 18.06.2024 Accepted : 25.07.2024)

## Abstract

The present investigation on “Studies on the effect of different combinations of soybean straw and sugarcane bagasse on yield of oyster mushroom” was conducted to evaluate different substrates combinations for growth and yield of oyster mushroom. The present study focused on evaluating the growth and yield of *Pleurotus* spp. using different substrate combinations of wheat straw, soybean straw and sugarcane bagasse which have economic approach in agro industry as the residues are readily available. Therefore, the aim of this study was to investigate the feasibility of different substrate combinations for cultivation of oyster mushroom. The experiments consist of six treatments with four replications in complete randomized design. Treatments of experiment are T<sub>1</sub> - Soybean straw, T<sub>2</sub> - Sugarcane bagasse, T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%), T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%), T<sub>5</sub> - Soybean straw (25%) + Sugarcane bagasse (75%), T<sub>6</sub> - Wheat straw. Average fruit weight varied from 5.93 to 8.34 g. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) recorded maximum average fruit body weight (8.34g) and treatment T<sub>2</sub> - Sugarcane bagasse (100) recorded the minimum average fruit weight (5.93 g). The number of fruits (per kg dry substrate) for first, second and third harvest varied from 49.16 to 97.41, 45.33 to 84.10 and 21.92 to 37.1 respectively. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) recorded maximum numbers of fruits per kg dry substrate for first and third harvest 97.41 and 37.1 fruits respectively while the treatment T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%) recorded maximum number of fruits per kg dry substrate (84.10 fruits) for second harvest. The total yield of fresh oyster mushroom varied from 711.6 to 1101.11g per kg dry substrate. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) gave maximum yield (1101.11 g kg<sup>-1</sup> dry substrate) and the treatment T<sub>2</sub> - Sugarcane bagasse (100%) recorded with the minimum yield (711.6 g kg<sup>-1</sup> dry substrate).

**Key words : Soybean straw, sugarcane bagasse, dry substrate, wheat straw, oyster mushroom.**

Mushroom belongs to the group of fungi and it has the ability to convert cellulosic and lignolytic plant wastes into palatable food. A toadstool or mushroom is a spore-bearing, fleshy plant organ of a fungus that grows above ground, on soil or on its food supply. Mushroom references are available in ancient literature like ‘Vedas’ and Bible’. There are different kinds of mushroom having various size, shape, colour, edibility and appearance. There are approximately 5000 different types of

mushrooms found all over the world. Out of this, more than 2000 species are edible and 300 belonging to 70 genera are reported from India. Among this few species have been brought under cultivation on commercial scale. Out of 200 edible species of prime mushroom, about 80 species have been grown experimentally, 20 cultivated commercially and 4 and 5 species produced on industrial level throughout the world (Chang and Miles, 1989). Mushrooms are known vegetarian’s meat due to the rich source of minerals and proteins. The proteins of mushroom are considered to be intermediate between that of vegetables and animals. Amino

1. Associate Professor, 2 Head, Department of Soil Science and Agril. Chemistry MPKV, Rahuri 413 722, 3 and 4. Sr. Res. Asstt.

acids required for different biological activities in human body were also found in oyster mushroom. Oyster mushroom productivity is maximum in a very short time providing more protein per unit area than any other crop. Due to the high content of nitrogen and protein, which increases the biological efficiency of oyster mushroom. Oyster mushroom also produces metabolites of medicinal and pharmacological value, such as antimicrobials, immune stimulants, antioxidants and antitumourals. *Pleurotus* is an efficient lignin degrading mushroom and can grow well on different types of substrates containing lignocellulosic materials. *Pleurotus* spp. can grow well in variable temperature conditions; hence they are ideally suited for cultivation throughout the year in tropical regions of the world. They are able to colonize and degrade a large variety of lignocellulosic residues and required very short time for the growth than other edible mushrooms (Singh *et al.*, 2019).

*Pleurotus* species' fresh fruiting bodies have a high moisture content (90.8%) and both fresh and dried oyster mushrooms are high in proteins (30.4%), fat (2.2%), carbohydrates (57.6%), fibre (8.7%) and ash (9.8%) with 345 K cal of energy per 100 g of dry weight. It also contains vitamins like thiamin (4.8 mg), riboflavin (4.7 mg) and niacin (1087 mg) as well as minerals (Pandey and Ghosh, 1996). Protein content of mushrooms varies from 4 to 44% according to the species (Crisan and Sands, 1978). *Pleurotus sajor-caju* is a commercially cultivated species of oyster mushroom. *P. sajor-caju* has high protein content and a low fat content. It also contains vitamins (B1, B2, C), minerals (P, Na, Ca), and a high content of fibres and carbohydrates. *Pleurotus sajor-caju* exhibited strong antioxidant properties. It also showed moderate antibacterial activity against *Streptococcus aureus* and *Vibrio cholerae*. The presence of vitamins and carotenoids in *P. sajor-caju* can

also play a protective role in diseases such as cancer and cardiovascular disease. The species is also known to possess antineoplastic (inhibiting or halting the development of a tumor, i.e., neoplasm) properties (Chandraprakash *et al.*, 2022).

Oyster mushrooms are the easiest, fastest and cheapest to grow, require less preparation time and production technology. Also, the first flush is usually large, without the need for compost, manure, limestone, casing or temperature shocks. With more than 100% biological efficiency, coupled with its distinctive flavour, aroma and excellent drying and preservation qualities, it is assured a unique status as a delicacy (Mandee *et al.*, 2005). Various agricultural by-products are being used as substrates for the cultivation of the oyster mushroom. Some of these wastes include banana leaves, peanut hull and corn leaves, mango fruits and seeds, sugarcane leaves, wheat and rice straw (Cangy and Peerally, 1995). Agricultural wastes such as maize cob, maize straw, wheat straw, palm kernel cake, cotton seed hull, saw dust, spent grain; grass families are good materials for farm substrate production. *Pleurotus* are efficient lignin degraders, which can grow on different agricultural wastes which makes *P. sajor-caju* cultivation as an excellent alternative for production of mushrooms when compared to other mushrooms (Kausar, 1998). In Europe, wheat straw is used, while in South East Asian countries sawdust is more common. The majority of these substrates can be used as animal feed. However, their low digestibility, low protein content and high lignin content render them unpopular and unacceptable (Mandee *et al.* 2005). The mushroom yield was related to chemical and biological composition of the substrate which is important for the growth of mushroom than other substrates (Tsegaye and Tefera 2017).

## Material and Methods

During the present research work on “Studies on the effect of different combinations of soybean straw and sugarcane bagasse on yield of oyster mushroom” a set of experiment was carried out at All India Coordinated Research Project on Mushroom, College of Agriculture, Pune. During present investigation the master spawn of *Pleurotus sajor-caju* was maintained and used from the AICRP on Mushroom, College of Agriculture, Pune. For bed preparation, high density polythene bags were used. For cultivation of oyster mushroom chopped dried soybean straw, sugarcane bagasse and wheat straw was used. Before bed filling, the fumigation of room was done by using Formaldehyde (37-41%) @ 2%. To avoid any harmful effect of formaldehyde, essential tools such as the well sanitized hand gloves and mask were used. For fumigation of oyster mushroom cultivation room, formaldehyde (37-41%) @ 2% was used. In addition, Bavistin and copper Sulphate were also used @ 10 % for fumigation of mushroom growing rooms. The bold seeded whole wheat grains were weighed and rinsed 2-3 times in tap water. The vase was filled with water until the grains were just immersed. The grains were boiled in hot water to prevent starch from escaping. After cooking, the grains were spread over a large wire gauze mesh to drain extra water. Overnight, the grains were maintained in a ventilated atmosphere. Based on the weight of these boiled grains, 4 percent calcium carbonate was mixed the next morning. Mother spawn was prepared in conical flasks, whereas commercial spawn was prepared in heat resistant polypropylene bags. The cooked grains were packed to 3/4 capacity in these conical flasks and polypropylene bags (size 20cm x 30cm gauge 175) and then plugged with non-absorbent cotton. The polypropylene bags and conical flasks were sterilized in an autoclave at 20 pounds PSI for 1 hr. After sterilization, the bags and flask were cooled and

thoroughly shaken. The flask and bags were placed in a sterile (UV-sterilized) laminar flow chamber. The experiments consist of six treatments with four replication in complete randomized design. T<sub>1</sub> - Soybean straw, T<sub>2</sub> - Sugarcane bagasse T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%), T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%), T<sub>5</sub> - Soybean straw (25%) + Sugarcane bagasse (75%), T<sub>6</sub> - Wheat straw. The watering was taken off for a day before harvesting. Harvesting of fruiting bodies was done before shedding of spores. The total number of fruiting bodies and fresh weight was noted immediately after harvesting.

## Results and Discussion

**1. Average fruit weight :** Results of the average fruit body weight are presented in Table 1. the average fruit body weight was determined by selecting 10 mature fruit bodies at random from each treatment. Average fruit weight varied from 5.93 to 8.34 g. The treatment T<sub>3</sub>- Soybean straw (75%) + Sugarcane bagasse (25%) recorded maximum average fruit body weight (8.34g) and treatment T<sub>2</sub> - Sugarcane

**Table 1.** Effect of different substrate combinations on average fruit body weight (g) of the oyster mushroom

Treat-ments	Average fruit wt. (g)
T <sub>1</sub> - Soybean straw (100%)	7.57 <sup>bc</sup>
T <sub>2</sub> - Sugarcane bagasse (100%)	5.93 <sup>a</sup>
T <sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%)	8.34 <sup>c</sup>
T <sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%)	6.99 <sup>abc</sup>
T <sub>5</sub> - Soybean straw (25%) + Sugarcane bagasse (75%)	6.55 <sup>ab</sup>
T <sub>6</sub> - Wheat straw (100%)	6.68 <sup>ab</sup>
SE(m) ±	0.50
C.D @ 5%	1.50

Note: Treatment means having common super subscripts are statistically non-significant. Treatment means having different super subscripts are statistically significant.

bagasse (100) recorded the minimum average fruit weight (5.93g). The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%), T<sub>1</sub> - Soybean straw (100%) and T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%) were found to be at par with each other. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) observed significantly superior to rest of the treatments.

The results are harmony with those of Deora et al. (2021), who found that *Pleurotus eryngii* had an average fruit body weight ranging from 25.3 to 96.4 g. In the experiments with *Pleurotus* spp., Sharma and Jandaik (1983) found that the fruit body weight ranged from 5.0 to 5.8 g. Hassan et al. (2010) found that the average fruit body weight in Peshawar ranged from 2.3 to 9.7 g. According to Khade et al. (2019), the average fruit body weight of *Hypsizygu* *ulmarius* ranged from 2.72 to 10.64 g.

**2. Number of fruits per kg dry substrate :** Results of the numbers of fruits per kg dry substrate at every harvest are presented in Table 2.

**2.1. First harvest :** Number of fruits per kg dry substrate at first harvest varied from 49.16 to 97.41. The maximum number of fruits (97.41) per kg dry substrate was found in T<sub>3</sub> -

Soybean straw (75%) + Sugarcane bagasse (25%) while the treatment T<sub>2</sub> - Sugarcane bagasse (100) recorded the minimum number of fruits (49.16) per kg dry substrate. T<sub>1</sub> - Soybean straw (100%) and T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) were found to be at par with each other. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) was found to be significantly superior over rest of treatments.

**2.2. Second harvest :** Number of fruits at second harvest varied from 45.33 to 84.10. The treatment T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%) observed with maximum number of fruits (84.10) per kg dry substrate and treatment T<sub>2</sub> - Sugarcane bagasse (100) recorded the minimum number of fruits (45.33) per kg dry substrate. The treatment T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%) reported maximum number of fruits than rest of treatments.

**2.3. Third harvesting :** Number of fruits for third harvest varied from 21.92 to 37.1 fruits per kg dry substrate. Maximum number of fruits (37.1) per kg dry substrate were found in T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) and the treatment T<sub>2</sub> - Sugarcane bagasse (100) recorded the minimum number of fruits (21.92) per kg dry substrate. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse

**Table 2.** Effect of different substrate combinations on number of fruits (per kg of dry substrate) after spwan run of oyster mushroom

Treat-ments	No. of fruits		
	First harvest	Second harvest	Third harvest
T <sub>1</sub> - Soybean straw (100%)	95.58 <sup>d</sup>	53.92	32.67 <sup>bc</sup>
T <sub>2</sub> - Sugarcane bagasse (100%)	49.16 <sup>a</sup>	45.33	21.92 <sup>a</sup>
T <sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%)	97.41 <sup>d</sup>	68.66	37.1 <sup>c</sup>
T <sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%)	78.91 <sup>c</sup>	84. 10	33.08 <sup>bc</sup>
T <sub>5</sub> - Soybean straw (25%) + Sugarcane bagasse (75%)	76.32 <sup>c</sup>	46.58	27.42 <sup>ab</sup>
T <sub>6</sub> - Wheat straw (100%)	60.75 <sup>b</sup>	54.46	26.42 <sup>ab</sup>
SE(m) ±	2.47	13.44	2.58
C.D @ 5%	7.38	NS	7.73

(25%), T<sub>1</sub> - Soybean straw (100%) and T<sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%) were statistically at par with each other.

The data obtained correspond with Khade et al. (2019) findings that *Hypsizygu* *ulmarius* generated 6.11 to 89.56 fruit bodies per bed. The findings are in conformity with those of Deora et al. (2021), who reported that *Pleurotus eryngii* produced 9.4 to 32.8 fruit bodies per bed. Gaikwad (2004) also reported that the total number of fruits produced by *Pleurotus sajor-caju* ranged from 125 to 200. *Pleurotus sajor-caju* produced a maximum of 65 fruit bodies per bed, reported by Shukla and Jaitley (2011).

**3. Yield of oyster mushroom :** Results of the yield performance are presented in Table 3. the yield of oyster mushroom varied from 711.60 to 1101.11g per kg dry substrate. The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) recorded maximum yield (1101.11g kg<sup>-1</sup> dry substrate) and the treatment T<sub>2</sub> - Sugarcane bagasse (100) recorded the minimum yield (711.60 g kg<sup>-1</sup> dry substrate). T<sub>1</sub> - Soybean straw (100%) and T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) were found to be at par with each other.

**Table 3.** Effect of different substrate combinations on yield of oyster mushroom

Treatment	Total yield (g kg <sup>-1</sup> dry substrate)
T <sub>1</sub> - Soybean straw (100%)	1039.54 <sup>c</sup>
T <sub>2</sub> - Sugarcane bagasse (100%)	711.60 <sup>a</sup>
T <sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%)	1101.11 <sup>c</sup>
T <sub>4</sub> - Soybean straw (50%) + Sugarcane bagasse (50%)	889.70 <sup>b</sup>
T <sub>5</sub> - Soybean straw (25%) + Sugarcane bagasse (75%)	821.67 <sup>ab</sup>
T <sub>6</sub> - Wheat straw (100%)	851.21 <sup>ab</sup>
SE(m) ±	49.44
C.D @ 5%	148.02

The treatment T<sub>3</sub> - Soybean straw (75%) + Sugarcane bagasse (25%) was found significantly superior to rest of treatments.

According to Dubey et al. (2019) the maximum yield (1515g) of oyster mushroom was recorded in paddy straw substrate followed by banana leaves (517.5g), wheat straw (480g) and sugarcane bagasse (98.75g). The findings correspond with those of Deora et al. (2021), who also reported that the yield of *P. eryngii* on wheat straw and rice straw was 767.4 g kg<sup>-1</sup> and 884.0 g kg<sup>-1</sup> dry substrate, respectively. Dhobale (2023) determined the yield of *P. sajor-caju* on wheat straw to 706.82 g kg<sup>-1</sup> to 1306.82 g kg<sup>-1</sup> dry straw. Sawant et al. (2022) measured the yield performance of *P. sajor-caju* on soybean straw, wheat straw, and sugarcane trash and found that it generated 752.64 g kg<sup>-1</sup>, 654.01 g kg<sup>-1</sup>, and 635.14 g kg<sup>-1</sup> dry substrate, respectively.

## References

- Cangy, C. and Peerally, A. 1995. Studies of *Pleurotus* production on sugar-cane bagasse. *African Journal of Mycology and Biotechnology.*, 3(2): 67-79.
- Chandraprakash, S., Kamalasureyan, C. M., Kumar, R. P., Manoj, K., Pradesh, G. J., Kumar, R. N. and Partheeban, C. 2022. Performance of Various Substrates in the Cultivation of oyster mushroom (*Pleurotus sajor-caju*). *Eco. Env. and Cons.*, 28: 252-256.
- Chang, S. T. and Miles, P. G. 1989. Edible mushrooms and their cultivation. CRC Press. Boca Raton, Florida. 6: 555-565.
- Crisan, E. V., and Sands, A. 1978. Nutritional value. In "The Biology and Cultivation of Edible Mushroom", eds. Chang, ST, and Hayes, New York: Academic Press: 137-168.
- Deora, A., Sharma, S. S. and Rohith, M. 2021. Cultivation of Kabul Dhingri (*Pleurotus eryngii*) mushroom by standardizing protocols in subtropical zones of world. *Scientific Reports*, 11: 14692.
- Dhobale, S. T. 2023. Evaluation of different lime formulation for substrate sterilization in oyster mushroom cultivation. A thesis submitted to M.P.K.V. Rahuri, Dist. Ahmednagar (MS), India

- Dubey, D., Dhakal, B., Dhami, K., Sapkota, P., Rana, M., Poudel, N. S. and Aryal, L. 2019. "Comparative study on effect of different substrates on yield performance of oyster mushroom," *Global Journal of Biology, Agriculture and Health Sciences*, 7(7): 1-7.
- Gaikwad, A. B., 2004. Effect of biofertilizers on yield of mushroom (*Pleurotus sajor caju*). A thesis submitted to M.P.K.V. Rahuri, Dist. Ahmednagar (MS), India.
- Hassan, S., Mohammad, A. and Khan, K. 2010. Cultivation of the oyster mushroom [*Pleurotus ostreatus* (jacq.) p. Kumm.] in two different agroecological zones of Pakistan. *African J. of Biotechnol.* Vol. 10(2): 183-188.
- Kausar, T. 1988. Cultivation of mushrooms using crop residues as substrate. Ph. D. Thesis. Department of Botany. University of Punjab. Lahore, Pakistan.
- Khade, R. S., Jadhav, A. C., Dhavale, M. C. and Gaikwad, A. P. 2019. Evaluation of Different Supplementations on Growth and Yield of Elm Oyster (*Hypsizygus ulmarius*) Mushroom. *Int. J. Curr. Microbiol. App. Sci.* 8(11): 1084-1095.
- Mandeel, Q. A., A-talth A .A., and Mohamed S .A. 2005. Cultivation of Oyster Mushrooms (*Pleurotus sp.*) on Various Lignocellulosics Wastes. *World J. Microbial. Biotechnol.* 21 (4): 601-607.
- Pandey, R. S. and Ghosh, S. K. 1996. A Handbook on Mushroom Cultivation, Emkay publ., Delhi pp: 134.
- Sawant, D. M., Pawar, R. M. and Purane, B. B. 2022. Cultivation of *Pleurotus sajor-caju* (F.) Singer on Different Agro Wastes. *Int. J. Creative Res. Thoughts*, 10: 2320-2882.
- Sharma, A. D. and Jandaik, C. L. 1983. Effect of spawn run duration on yield and some quality parameters of oyster mushroom (*P. sajor-caju*). *Indian J. Mushrooms*, IX :7-11.
- Shukla, S. and Jaitley, A. K. 2011. Morphological and biochemical characterization of different oyster mushroom (*Pleurotus spp.*) *J. of Phytol.* 3(8): 18-20.
- Singh, U. P., Maurya, K., and Singh, D. 2019. Evaluation of agro-wastes for the production of oyster mushroom (*Pleurotus sajor caju*). *Journal of Pharmacognosy and Phytochemistry*, 8 (3): 231-234.
- Tsegaye Z, and G. Tefera, 2017. "Cultivation of oyster mushroom (*Pleurotus ostreatus* kumm, 1871) using agro-industrial residues," *Journal of Applied Microbiological Research*, 1 (1): 1-6.
-