

Lime and Manure Application to Low-fertility Tropical Soils Enhances Phosphorus Bioavailability for Increased Agronomic Productivity

V. U. Ugwu, A. I. Orah, C. I. Osuji, J. C. Akubue, S. E.e Obalum*, C. A. Igwe

Dept. of Soil Science, Faculty of Agriculture, University of Nigeria, Nsukka 410001, Enugu State, Nigeria

*Corresponding author's Email : sunday.obalum@unn.edu.ng;

(Received : 24.07.2024 Accepted : 22.11.2024)

Abstract

Liming and manuring to ameliorate soil acidity and enhance mineralization in acid tropical soils could promote their agronomic productivity. Synthetic lime and poultry-droppings manure's effects in sandy-loam Ultisols were evaluated on soybean growth, exploring its relationships with soil pH, soil available P and plant P content. Lime was applied at 0, 2.5 and 5 t ha⁻¹ equivalents and manure at 0, 25 and 50 t ha⁻¹ equivalents to 2.5-kg potted soils. Crop growth data were collected during and soil-plant analysis after six weeks of growth. Treatment had more pronounced effects on plant height than leafiness, being generally highest in lime 2.5 + manure 50 where the plants were about 3 x taller compared to unamended control. Soybean grew better due to 2.5 than 0 and 5 t ha⁻¹ lime, and to 50 than 0 and 25 t ha⁻¹ manure. Plant height and leaf area depended on soil pH's influence on soil available P ($R^2 = 0.69^{**}$) and plant P content ($R^2 = 0.85^{**}$), respectively, while above-soil biomass on soil pH or plant height ($R^2 = 0.74^{**}$). Moderate liming (2.5 t ha⁻¹) with ample manuring (50 t/ha) could enhance crop early-stage vegetative growth in low-fertility tropical soils, due largely to amelioration of soil acidity to enhance uptake of bioavailable P.

Key words : Acid tropical soils; synthetic lime; nutrient-rich manure; soil available P; plant nutrition.

Highly weathered tropical soils are known to be naturally acidic and of low fertility status (Caires *et al.*, 2008; Fageria *et al.*, 2011; Ubi *et al.*, 2017). These acidic soils with high levels of rainfall over years are leached of basic ions (Ubi *et al.*, 2017). Heavy rainfall is thus one major cause of basic cations removal through leaching over a long period of time. Leaching increases acidity as leached soils are left with toxic and insoluble compounds of iron (Fe) and aluminium (Al) in soil (Zhang *et al.*, 2016). Acidic soils, therefore, have high amount of Fe and Al oxides with high specific surface area (Achatet *et al.*, 2010). These oxides have high appetite for phosphate which can sometimes become almost insatiable (Bueis *et al.*, 2019). Oxides of Fe and Al are the main sorbents of phosphate responsible for phosphorus (P) sorption and thus its unavailability to plants in acidic soils (Turrión *et al.*, 2008; Achatet *et al.*, 2010; Bueis *et al.*, 2019). Poor availability of P

is a major limiting constraint for crop production in acidic soils (Wang *et al.*, 2010).

Tropical soils are also of low fertility with low soil organic matter (SOM) content due to high mineralization rate. The SOM influences the physical, chemical, and biological characteristics of the soil (Adiaha, 2017; Sandirakirana and Arifin, 2021). Depletion of SOM adversely affects P bioavailability and thus limits crop productivity in these acid low-fertility soils. By contrast, increases in SOM imply greater ability of soil to resist the natural tendency of becoming acidic (Ejersa, 2021). Soil content of SOM influences the activities of soil microbes which transform organic P into inorganic P (Allison and Vitousek, 2005; Bueis *et al.*, 2019), and its decomposition releases inorganic P, a form accessible by plants (Gichangi, 2019). Organic acids originating from SOM decomposition enhance the solubility of calcium (Ca)

phosphates, and these organics are adsorbed by Fe and Al oxides, blocking P sorption sites (Bueis *et al.*, 2019). Also, SOM is a major constituent of soil sorption complex which is responsible for the binding of ions in the soil and thus plays a critical role in P sorption (Debicka *et al.*, 2016; Chukwuma *et al.*, 2024).

Liming is a common management practice in most agricultural systems used to mitigate soil acidification toward enhancing crop productivity in such soils (Mkhonza *et al.*, 2020). The application of lime brings about desirable soil pH, decreases toxicity of Fe and Al, increases Ca and magnesium supplies as well as enhances the availability of P (Mkhonza *et al.*, 2020). Increased soil pH, resulting from lime application, increases microbial activity (Rousket *et al.*, 2020; Paradelo *et al.*, 2015). This in turn enhances decomposition of SOM (Garbui *et al.*, 2011; Grover *et al.*, 2017). Lime application increases the availability of nutrients, which would otherwise be strongly limited by low soil pH (Rastija *et al.*, 2014), improving nutrient uptake (Takala, 2019; Mkhonza *et al.*, 2020). Lime application increases P availability by decreasing the amount of Fe and Al ions and P fixation on their oxides (Mkhonza *et al.*, 2020). Liming creates a favourable root environment for plants as a result of these different effects it has on soils.

Manures can be a valuable, relatively cheap source of P and other essential plant nutrients (Han *et al.*, 2008; Saleem *et al.*, 2017; Obalum *et al.*, 2020). Mineralization following manure addition to soils enhances nutrient availability by transforming nutrient elements from organic to inorganic forms (Saleem *et al.*, 2017). Application of manures to acid soils can boost SOM which being negatively charged complexes Fe and Al ions, inhibiting their crystallization and role in P sorption and/or precipitation as insoluble phosphates in soils (Haynes and Mokolobate, 2001; Brady and

Weil, 2008). Through adsorption site competition, organic acids in manures also inhibit phosphate adsorption (Borggaard *et al.*, 2005). Soil application of manures thus promote arable crops production via increased P bioavailability (Haynes and Mokolobate, 2001; Adiaha, 2017).

Synthetic limes can be combined with P fertilizers for increased crop productivity in tropical Africa (Desalegn *et al.*, 2016; Amsalu and Beyene, 2020; Tshiabukole *et al.*, 2022). However, combining lime and organic P source in the coarse-textured soils of the region might, by alleviating the toxic effects of Fe-Al oxides (Islam *et al.*, 2021), ensure non-sorption of the released P. Poultry droppings can have more liming and P-supplying effects in acid tropical soils than other manures of organic origin (Adubasime *et al.*, 2018; Ugwu *et al.*, 2020; Chukwuma *et al.*, 2024). In such soils, a compatibility question mark is hanging on co-application of ash as organic lime and poultry-droppings manure (Nwite *et al.*, 2013), but impressive agronomic responses had been found for co-application of synthetic lime and this manure (Nnadi *et al.*, 2020). Lime-manure mix not only aligns with the global principle of organic-inorganic complementation in soil and water management, but is also more practicable than co-application of ash and rock phosphate that is among the P-fertility management options proposed for the soils (Ndzeshala *et al.*, 2021).

In the present study, we opted to combine lime and poultry-droppings manure, justified by the greater ability of lime to raise soil pH compared to ash (Materechera and Mkhabela, 2002) and the high cost of ash equivalent of lime for the variable-charge and pH-buffered soil investigated (Ndzeshala *et al.*, 2021), vis-a-vis this manure's prized effectiveness. Research on the synergy of these soil amendments in the tropics has, however,

used rather low application rates of especially the manure component (Islam *et al.*, 2022; Kabango *et al.*, 2022; Akpan *et al.*, 2023). Therefore, objectives of this study were to evaluate the agronomic responses to the effects of co-application of synthetic lime and poultry-droppings manure at standardized rates of the former in coarse-textured tropical soils, determine their effects on P availability in the soils, and examine the relationships between crop growth and soil available P and plant uptake of it.

Materials and methods

Soil of the study : The research was conducted using soil from the University of Nigeria Teaching and Research Farm, which was collected some 5 m away from the glasshouse of the Department of Soil Science (06°51'47" N, 07°25'22" E; 443 m asl) at Nsukka campus of the University. The climate is humid tropical, with mean annual rainfall of 1600 mm, mean minimum/maximum temperatures of 21/31 °C, and relative humidity in the range of 55-90%. Soils of the area are mostly underlain by false-bedded sandstone or upper-coal measure. At the surface, they can be excessively 'porous' and hence well-drained and often prone to drought (Obalum and Obi, 2014). The occurrence of the soils in a high-rainfall zone, coupled with their 'porous' attribute, makes them highly leached of basic cations and hence strongly acid (Obalum *et al.*, 2011a). By the prevailing climate/soil-supported cum anthropogenic vegetation, the area is a Derived Savannah in southeastern Nigeria.

The soil used for the study, derived from false-bedded sandstone, is deeply weathered and of oxide-rich mineralogy. It is hence characterized by coarse texture and reddish brown moist colour, belonging to the order Ultisol of the USDA's Soil Taxonomy. The particle size distribution of the surface horizon

(sand, silt and clay contents of 750, 70 and 180 g kg⁻¹, respectively) places the soil in the texture class of Sandy Loam and qualifies it as a coarse-textured one.

Procurement of study materials :

Topsoil (0-20 cm) was collected from a part of the Teaching and Research Farm that had been under fallow for over three years. The heap of soil was spread and left to air-dry in the glasshouse for about two weeks. Thereafter, it was crushed and sieved using 2-mm sieve to remove any gravels and dry plant parts before potting. Before the pot trials, the air-dry topsoil showed acidic soil pH of 5.2, soil organic carbon and total nitrogen contents of 13.78 and 0.53 g kg⁻¹, respectively, available P content of 5.20 mg kg⁻¹, and apparent cation exchange capacity of 13.60 cmol kg⁻¹.

Quicklime (CaO) with calcium carbonate equivalent of 88% (CaO-88%) and treated seeds of a soybean variety with an indeterminate growth habit used as the test crop were bought from the local market at Nsukka, southeastern Nigeria. Poultry droppings were sourced from the Battery Cage System of the Animal Science Section of the Farm. Curing of the fresh poultry-droppings manure was done sun-drying for one week. Then, it was crushed and sieved to allow for proper mixing with the soil. A proximate analysis on the dry manure showed that it had a pH of 8.8 and C, N, P, K, Ca, Mg and Na contents of 283, 24.50, 5.20, 6.50, 11.00, 4.90 and 2.40 g kg⁻¹, respectively.

Experimental design and Procedure :

The research was executed as a glasshouse study, using ceramic pots of approximate capacity 2.25 L with drainage hole at the bottom. Treatments consisted of factorial combination of three rates (0, 2.5 and 5.0 t ha⁻¹) of quicklime and three rates (0, 25 and 50 t ha⁻¹) of poultry-droppings manure. Lime rates were adopted after 2.5 t ha⁻¹ for similar sandy-

loam Ultisols (Agba *et al.*, 2017), while manure rates were adopted after the optimum 'agronomic' rate for the sandy-loam Ultisols investigated, found to be 20-25 t ha⁻¹ (Ogunezi *et al.*, 2019; Onah *et al.*, 2023). The factorial combination gave nine treatments replicated three times in a completely randomized design (CRD). These treatments were mixed with 2.5 kg air-dry topsoil and, thereafter, placed into the pots. Assuming a soil bulk density of 1,500 kg m⁻³, the application rates translated into 0, 2.5 and 5.0 g per potted soil for quicklime and 0, 25 and 50 g per potted soil for manure. These soil addition rates are represented as L₀, L_{2.5} and L_{5.0} for lime rates and PD₀, PD₂₅ and PD₅₀ for manure rates. The three replications of these nine treatments gave a total of 27 potted soils for the study.

The potted soils were watered to field capacity and left overnight before sowing soybean. Two seeds were sown per potted soil. One week later, the young seedlings were thinned to one stand per potted soil. From soybean sowing till six weeks after sowing (WAS) when the pot experiment was terminated, the potted soils/seedlings were watered to pre-determined field capacity at two-day intervals, while manually removing weeds as they emerged. The glasshouse had its side louvers wide open such that the environmental conditions including lighting duration, temperature, relative humidity, etc. in and outside the glasshouse were the same throughout the study.

Agronomic data collection : Plant morphological growth data were collected weekly during 2-6 WAS. Growth parameters considered were plant height (from soil level to the tip of plant); leaf area, estimated as 0.31LW (with L and W as average lengths and widths of two broadest leaves of the seedling, respectively); and number of leaves per plant. At 6 WAS, the plants were cut off from their soil

base and allowed to undergo curing at the prevailing ambient temperature at the glasshouse for 48 h before weighing to obtain the above-soil biomass. Plant leaf samples were collected from the cured biomass for P content analysis. This aligned with the proposition of 6 WAS of soybean as its vegetative growth stage for plant nutrient analysis on the mature leaves (Stammer and Mallarino, 2018). The leaf samples were air-dried further for three weeks before the analysis.

Laboratory analyses : Soil samples were air-dried to constant weight, crushed and passed through a 2-mm mesh sieve before analyses. Soil pH was determined in suspensions of soil in deionized water in a solid-liquid ratio of 1:2.5 (McLean, 1982), and the values measured potentiometrically using a Beckman's zero-matic glass electrode pH metre. Available P was extracted with Bray 2 solution; thereafter, the values were measured colorimetrically using the method described by Olsen and Sommers (1982). Plant P content was analyzed by digesting 0.5 g of plant samples with 20 ml each of nitric acid and perchloric acid, after which the digest was washed with distilled water and later extraction was done using Bray 2 solution (Olsen and Sommers, 1982).

Data analyses : Data were analysed using GenStat and SPSS software. First, two-way analysis of variance was used to test for differences in treatment effects using GenStat. Where significant differences existed, the means were separated by the Fisher's least significant difference procedure at $p = 0.05$ (F-LSD 0.05). Also, Pearson's bivariate correlations and stepwise multiple linear regressions were carried out using SPSS. This was to examine the relationships among the vegetative growth indices of soybean, soil pH, soil content of available P, and soybean plant P content.

Results and Discussion

Effects of lime and manure rates on soybean growth parameters : The effects of lime and manure rates on plant height, leaf area and number of leaves per soybean plant are shown (Figures 1, 2 and 3, respectively). Generally, plant height differed among treatments. The data show a trend of highest values in treatment $L_{2.5} + PD_{50}$. Leaf area, however, differed among treatments only at 3 and 4 WAS. The trend observed in leaf area was same as for plant height; highest values were consistently observed in treatment $L_{2.5} + PD_{50}$. Number of leaves differed among treatments at 4, 5 and 6 WAS. The highest values for this growth parameter were observed in treatment $L_{2.5} + PD_{50}$ at 2 and 3 WAS, while treatment $L_{2.5} + PD_{25}$ showed highest values at the later growth stages being 4, 5 and 6 WAS. For all growth parameters measured in this study, lowest values were generally observed in treatment $L_0 + PD_0$ (control).

The main effects of treatment showed that lime application rate affected soybean growth. With the exception of plant height trend of $L_5 > L_{2.5} > L_0$ during 2 WAS, plant height and leaf area generally differed thus $L_{2.5} = L_5 > L_0$ during 2 and 5 WAS, but $L_{2.5} > L_5 > L_0$ and $L_{2.5} > L_5 = L_0$, respectively during 3, 4 and 6 WAS. So, apart from 2 WAS, lime rate of 2.5 t ha^{-1} produced the tallest plants all through the growth phase of the soybean plants. For number of leaves per plant, the trend was $L_{2.5} = L_5 > L_0$ during 2 and 3 WAS, but $L_{2.5} > L_5 > L_0$ during 4, 5 and 6 WAS. The effects of poultry-droppings manure were rather more definite in that all three growth indices of soybean generally differed thus $PD_{50} > PD_{25} = PD_0$ during 2 and 3 WAS and $PD_{50} > PD_{25} > PD_0$ during 4, 5 and 6 WAS. Overall, liming at 2.5 t ha^{-1} , on one hand, and applying manure at 50 t ha^{-1} , on the other hand, enhanced the vegetative growth of soybean the most.




Fig. 1. Effects of synthetic lime and poultry-droppings manure application to the coarse-textured acid soil on plant height (cm) of the soybean variety with an indeterminate growth habit used as test crop ($n = 27$).

L_0 , $L_{2.5}$ and $L_{5.0}$ - lime at 0, 2.5 and 5 t ha^{-1} , respectively; PD_0 , PD_{25} and PD_{50} - poultry-droppings manure at 0, 25 and 50 t ha^{-1} , respectively; WAS-weeks after sowing Error bars represent Fisher's least significant difference at $p \leq 0.05$ (F-LSD $_{0.05}$).




Fig. 2. Effects of synthetic lime and poultry-droppings manure application to the coarse-textured acid soil on mean leaf area (cm^2) of the soybean plant ($n= 27$).

L_0 , $L_{2.5}$ and $L_{5.0}$ - lime at 0, 2.5 and 5 t ha^{-1} , respectively; PD_0 , PD_{25} and PD_{50} - poultry-droppings manure at 0, 25 and 50 t ha^{-1} , respectively; WAS-weeks after sowing; NS - not significant at $p \leq 0.05$ Error bars represent Fisher's least significant difference at $p \leq 0.05$ (F-LSD $_{0.05}$).




Fig. 3. Effects of synthetic lime and poultry-droppings manure application to the coarse-textured acid soil on number of leaves per soybean plant ($n = 27$).

L_0 , $L_{2.5}$ and $L_{5.0}$ —lime at 0, 2.5 and 5 $t\ ha^{-1}$, respectively; PD_0 , PD_{25} and PD_{50} —poultry-droppings manure at 0, 25 and 50 $t\ ha^{-1}$, respectively; WAS – weeks after sowing; NS—not significant at $p \leq 0.05$ Error bars represent Fisher's least significant difference at $p \leq 0.05$ (F-LSD_{0.05}).

There was a general trend of treatments with lime and manure promoting soybean growth more than sole application of either. Thus, liming and manuring could have synergistic effects in plant growth in acid low-fertility soils. Liming creates a favourable root environment by ensuring desirable soil pH, decreasing Fe and Al toxicity, and enhancing the availability macronutrients. It also improves soil biological activity and mineralization of organic compounds, thereby improving nutrient uptake (Saleem *et al.*, 2017; Takala, 2019). Also, the addition of manures increases microbial activity that ensures the biochemical transformations and mineralization of plant nutrients (Saleem *et al.*, 2017). Liming ensured nutrient elements made available from the manure were accessible for plant uptake due to its effect of decreased acidity. Appropriate combination of synthetic lime and mineral fertilizers could improve the bioavailability of plant nutrients for enhanced crop growth and yields (Ameyu, 2019; Nnadi *et al.*, 2020).

Treatment $L_{2.5} + PD_{50}$ tended to show the most pronounced effect on soybean vegetative growth. The optimum 'agronomic' rate of poultry manure for the sandy-loam soil investigated is 20-25 $t\ ha^{-1}$ (Ogunziet *et al.*, 2019; Onah *et al.*, 2023). So, the $L_{2.5} + PD_{50}$ represents co-application of synthetic limes at moderate rates of $\leq 2.5\ t\ ha^{-1}$ and poultry-droppings manure at $\leq$ double the subsisting 'agronomic' rate. Treatment $L_{2.5} + PD_{50}$ outperforming $L_{5.0} + PD_{50}$ indicates the negative effect of over liming on plant growth. Coarse-textured acid tropical soils may thus require low application rates of synthetic limes in the presence of high rates of effective animal manures.

Treatment effects on above-soil biomass, soil pH and available P, and plant P content :

At the end of six-week growth phase, above-soil biomass of the soybean plant as well as soil pH, available P and plant P content were assessed (Table 1). Lime effects on soybean above-soil biomass 6 WAS were such that treatments $L_0 + PD_0$ and $L_{2.5} + PD_0$ were similar and lower than $L_5 + PD_0$, unlike poultry-droppings manure effect that showed significant increases with application rate. Treatments $L_0 + PD_{25}$ and $L_{2.5} + PD_{25}$ were similar unlike $L_0 + PD_{50}$ that produced lower than $L_{2.5} + PD_{50}$ and $L_5 + PD_{50}$ for which values were similar. These results imply that synthetic lime may not be necessary with poultry-droppings manure at the so-called 'agronomic' rate, but at about the double of this application rate when the lime also needs not be applied beyond the moderate rate of 2.5 $t\ ha^{-1}$. Notably, merging $L_0 + PD_{50}$ and the control-like $L_{2.5} + PD_0$ gave rise to $L_{2.5} + PD_{50}$, while merging $L_0 + PD_{50}$ and $L_5 + PD_0$ both of which were similar gave $L_5 + PD_{50}$. From the standpoint of both the economy of agronomic production and the synergy of lime and manure over sole use of either, therefore, $L_{2.5} + PD_{50}$

was the optimal lime-manure combination in this study.

Where the adoption of $L_{2.5} + PD_{50}$ is cost-inhibitive, the resource-poor farmer can opt for the runner-up treatment being $L_0 + PD_{50}$ or $L_5 + PD_{25}$, whichever is more viable. Halving manure and adding CaO lime which the second of these two treatments is to the first has similarly been reported to be equivalents in soybean production in low-fertility tropical soils (Verde *et al.*, 2013). The next in performance was $L_5 + PD_0$ and it was similar to $L_0 + PD_{25}$.

Treatments $L_{2.5} + PD_{50}$ and $L_{5.0} + PD_{50}$ being similar in plant above-soil biomass are attributed to the low clay content of the soil and the overall environmental setting of the study, lime and manure types used and their application rates tested, and/or the short-duration nature of the study (Moreira *et al.*, 2015; Kabango *et al.*, 2022; Akpan *et al.*, 2023). The present observation indicates that CaO lime is required at $\leq 2.5 \text{ t ha}^{-1}$ in the presence of a P source for growing arable crops in acid loamy soils of the humid tropics (Opala, 2017). Beyond this rather moderate rate of lime, the corresponding rate of manure (about

twice the 'agronomic' rate) needed to maintain production at optimal levels would initially not change, after which an inverse lime-manure rate relationship sets in. From the study site, Nnadi *et al.* (2020) reported that co-application of CaO lime at 10 t ha^{-1} and poultry-droppings manure at 20 t ha^{-1} increased sweet potato yields. By its liming/SOM-enhancing effects in this coarse-textured soil, $L_{2.5} + PD_{50}$ would improve soil fertility (Umeugokweet *et al.*, 2021), soil aggregation/structure (Obalum *et al.*, 2024), or both (Ogunziet *et al.*, 2019; Obalum *et al.*, 2020; Ogumba *et al.*, 2024), leading to increased crop vegetative growth (Ebido *et al.*, 2021). For soybean, however, any improvements in soil aggregation/structure most unlikely contributed to the highest vegetative growth due to this treatment (Obalum *et al.*, 2011b; Osakwe *et al.*, 2023).

Treatment affected soil pH, available P and plant P content. Soil pH was highest and lowest in $L_5 + PD_{50}$ and $L_0 + PD_0$ (control), respectively, with generally higher values with (7.0-7.4) than without (6.6-7.1) combination of lime and manure (Table 1). Soil available P and plant P content were highest in $L_{2.5} + PD_{50}$, followed by $L_0 + PD_{50}$, while the lowest values

Table 1. Treatment effects on above-soil biomass of the indeterminate soybean plant, soil pH, soil available P, and plant P content, all assessed at six weeks of age of the plants ($n = 27$)

Treatment	Above-soil biomass (g 2.5^{-1} -kg-potted-soil)	Soil pH _{water}	AvP (mg kg^{-1})	% Plant-P
$L_{5.0} + PD_{50}$	18.50	7.4	27.98	0.28
$L_{5.0} + PD_{25}$	14.20	7.2	27.98	0.19
$L_{5.0} + PD_0$	11.87	6.7	9.33	0.15
$L_{2.5} + PD_{50}$	17.19	7.3	48.50	0.33
$L_{2.5} + PD_{25}$	11.40	7.0	22.38	0.16
$L_{2.5} + PD_0$	9.63	6.6	7.46	0.14
$L_0 + PD_{50}$	13.00	7.1	29.85	0.32
$L_0 + PD_{25}$	10.97	6.8	20.52	0.16
$L_0 + PD_0$ (Control)	8.27	5.8	5.60	0.12
F-LSD (0.05)	1.47	0.06	0.16	0.01

L_0 , $L_{2.5}$ and $L_{5.0}$ - lime at 0, 2.5 and 5 t ha^{-1} , respectively; PD_0 , PD_{25} and PD_{50} - poultry-droppings manure at 0, 25 and 50 t ha^{-1} , respectively; AvP - available phosphorus; Plant-P - plant P content

were from the control. Liming indirectly influences mineralization of manure-added nutrients by increasing soil pH which creates favourable conditions for microbial activities (Dada and Ewulo, 2011), and this leads to increases in plant uptake of the nutrients (Islam *et al.*, 2021). Notably, treatments $L_5 + PD_0$ and $L_0 + PD_{25}$ bearing only lime and only manure, respectively were similar in above-soil biomass, soil pH and plant P content but differed widely in soil content of available P. By this observation, 5 times the application rate of CaO lime would be needed of poultry-droppings manure to produce a similar effect on soil pH, implying that this manure's equivalent of CaO lime for the soil studied is 20%. Again, the observation indicates dependence of agronomic response to lime-manure amendments on soil pH-regulated bioavailable P rather than soil available P.

The main effects of treatment showed that both lime and manure contributed to the variations in the four plant and soil variables just presented. Soybean above-soil biomass and soil pH decreased with lime application rate ($L_{5,0} > L_{2,5} > L_0$), unlike soil available P ($L_{2,5} > L_5 > L_0$) and plant P content ($L_{5,0} = L_{2,5} > L_0$); while all four variables decreased with manure rate ($PD_{50} > PD_{25} > PD_0$). The increases in soil pH with an increase in lime rate can be associated with a decrease in hydrogen ions by its neutralization effect and an increase in hydroxyl ions following application of lime (Bolan *et al.*, 2003; Antoniadis *et al.*, 2015; Li *et al.*, 2019). Lime effect on soil available P suggests a lime-induced precipitation of Fe and Al as insoluble compounds (Antoniadis *et al.*, 2015) and hence decrease of Fe and Al binding/fixation of P at the optimal rate of lime, beyond which Ca fixation of P occurs (Amsalu and Beyene, 2020). Also, the deprotonation of Fe and Al hydroxides and the accompanying increases in negative surface charge with increasing soil pH contributes to an increase in available P (Opala, 2017). The higher Plant P content in limed potted soils

could be explained by enhanced release of soil nutrients including P, making it available for plant uptake.

Increasing soil pH with increase in manure rate in this study could be attributed to the release of cations following the mineralization of SOM. The increased presence of cations counters acidity and decreases exchangeable Al in soils amended with manure. Addition of manure has severally been reported to increase soil pH (Soremiet *et al.*, 2017; Ogunezi *et al.*, 2019; Okebalama *et al.*, 2020; Ndzeshala *et al.*, 2023; Onah *et al.*, 2023; Chukwuma *et al.*, 2024). Increases in P availability with manure rate can be attributed to organic P mineralization in the manure. Organic acids from SOM adhering to sorbing surfaces can inhibit phosphate adsorption via adsorption site competition, whereby they mask and prevent fixation sites from interacting with phosphate ions in solution (Borggaard *et al.*, 2005; Brady and Weil, 2008). Also, these negatively charged organic molecules adsorb or complex Fe-Al ions in acid soils (Haynes and Mokolobate, 2001; Brady and Weil, 2008). This could inhibit their crystallization and decrease their role in P sorption or precipitation. The phenomenon of increases in soil available P following manure application has been severally reported in the study area (Ogunezi *et al.*, 2019; Obalum *et al.*, 2020; Okebalama *et al.*, 2020; Umeugokwe *et al.*, 2021; Ndzeshala *et al.*, 2023; Onah *et al.*, 2023; Chukwuma *et al.*, 2024).

The increases in plant P uptake and content due to poultry-droppings manure addition are attributed to increased availability of P as a result of organic matter mineralization. Manure is known to be a reservoir of essential macro-nutrients and the nutrients are released upon decomposition and mineralization (Ogunezi *et al.*, 2019; Nnadi *et al.*, 2021). Improved availability of soil nutrient contents following poultry manure addition was reported to

increase uptake and hence concentrations of nutrients in plant tissues (Ewuloet *al.*, 2008; Ogunezi *et al.*, 2019).

Relationships among soybean growth, soil pH, soil available P, and plant P content : The coefficients of the bivariate correlations among growth/productivity indices of soybean, soil pH in water, soil available P, and plant P content are presented (Table 2). There were positive correlations among growth/productivity indices, all of which were significant except that between above-soil biomass and number of leaves. Soil pH, soil available P and soybean plant P content all had significant positive correlations with plant height, leaf area and above-soil biomass, but non-significant positive correlation with number of leaves. Also, soil pH correlated positively with soil available P and plant P content, both of which were also positively correlated. Soil pH often shows weak correlations with available P in the absence of soil acidity-ameliorating amendments in Nigeria (Obalum *et al.*, 2012; Obalum and Chibuike, 2017; Ukabiala *et al.*, 2021). Therefore, soil pH correlating with soil available P here could be attributed to the lime-manurial amendments' alteration of the former (Sato and Comerford, 2005; Ogunezi *et al.*, 2019; Chukwuma *et al.*, 2024) and/or to the use of one soil type of similar texture (Ifeanyi-Onyishi *et al.*, 2024). As

found here, P extractable from soil can be directly related to P concentrations in plants (Bolan *et al.*, 2003).

Soybean growth/productivity indices were regressed on soil pH, soil available P and plant P content (Table 3). Plant height was predicted by soil available P and leaf area by plant P content. Number of leaves per plant was not predicted any of the three independent variables, understandably because it was the only plant growth index that had no correlations with these indices. Instead, it was influenced by plant height. Soybean above-soil biomass was predicted by soil pH. However, repeating this regression analysis with plant height, leaf area and number of leaves included among the independent variables showed the soil available P-controlled plant height to be an alternative, equal-strength predictor of above-soil biomass. This points not just to the inter-relationships among plant height, above-soil biomass, soil pH and soil available P, but also to the underlying influence of soil pH on available P's control of biomass production in coarse-textured, low-fertility soils of the humid tropics.

The dependence of soybean growth/productivity on soil pH, available P, and plant P content was expected because of treatment-mediated influence of soil pH on P availability in

Table 2. Matrix of the coefficients of correlations (r) among growth/productivity indices of soybean†, soil pH in water, soil available P and plant P content (n = 9)

	Plant height	Leaf area	No. of leaves	Above-soil biomass	Soil pH _{water}	Soil AvP	Plant-P
Plant height	-						
Leaf area	0.85**	-					
No. of leaves	0.83**	0.74*	-				
Above-soil biomass	0.86**	0.85**	0.57	-			
Soil pH _{water}	0.82**	0.76*	0.66	0.86**	-		
Soil AvP	0.83**	0.90**	0.66	0.81**	0.80*	-	
Plant-P	0.72*	0.92**	0.59	0.81**	0.72*	0.86**	-

†For plant height, leaf area and number of leaves, mean values for the five consecutive sampling times (2, 3, 4, 5 and 6 weeks after sowing) were used for the correlations. AvP – available phosphorus; Plant-P – plant P content; * and **significant at p 0.05 and 0.01, respectively

the soil studied (Ogunezi *et al.*, 2019; Chukwuma *et al.*, 2024), especially with its not varying in soil texture (Ifeanyi-Onyishi *et al.*, 2024). It was also because of soil pH's underlying influence on plant's preferential uptake of P from this soil for increased crop productivity (Ndzeshala *et al.*, 2023). Similar to our data, Ndzeshala *et al.* (2023) and Ugwu *et al.* (2024) reported that about 88% of the variations in maize growth responses to nutrient-supplying organic amendments in potted coarse-textured, low-fertility soils of the study area to be in line with soil available P content. Treatment $L_{2.5} + PD_{50}$ being the optimal lime-manure combination, lime main effect on soil available P being $L_{2.5} > L_5 > L_0$, and soybean P nutrition playing a prominent role in its productivity are in unison.

With the role of soybean P nutrition in its biomass productivity in this study highlighted, the major mechanism behind $L_{2.5} + PD_{50}$'s effectiveness was most likely decreased Fe-Al fixation of P with increasing soil pH (Amsalu and Beyene, 2020). This $L_{2.5} + PD_{50}$ could not outperform $L_5 + PD_{50}$, suggesting that the manure component of these treatments, being above the optimum 'agronomic' rate of 20-25 t ha⁻¹ for this soil (Ogunezi *et al.*, 2019; Onah *et al.*, 2023), supplied P in excess of soybean requirement. So, the liming above the optimal rate of 2.5 t ha⁻¹ in treatment $L_5 + PD_{50}$ probably led to Ca fixation of the excess P due

to this treatment (Amsalu and Beyene, 2020), such that it would not contribute to further P nutrition and hence biomass accumulation by the plant.

Therefore, $L_{2.5} + PD_{50}$ is adopted as the optimal lime-manure combination for arable crops in the study area and similar humid tropical agro-environments. The lime-manure ratio in $L_{2.5} + PD_{50}$ is 0.05. This is rather low compared to the most effective ratio of 0.13 from a field-based evaluation of soybean responses to dolomite lime and cattle-dung manure effects in a clayey Brazilian Oxisol (Moreira *et al.*, 2015). The use of potted soils in our study and differences in environmental setting, soil type, and lime-manure types may explain the lower ratio in the adopted $L_{2.5} + PD_{50}$.

Conclusion

As a test crop, soybean showed interesting agronomic responses to CaO lime and/or poultry-droppings manure effects in the acid sandy-loam soil studied. Co-application of these synthetic and organic amendments at the rates of 2.5 and 50 t ha⁻¹, respectively to the soil enhanced the crop's vegetative growth and above-soil biomass better than the other rate-varying options of lime-manure combinations. The increases in soybean productivity due to this

Table 3. Results of multiple linear regressions of growth/productivity indices of soybean† on soil pH, soil available P and plant P content (n = 9)

Regression models	R ²	Adj. R ²	SEE
Plant height = 39.986 + 1.502AvP	0.692	0.648	14.559
Leaf area = 2.037 + 55.062plant-P	0.852	0.831	1.998
Number of leaves = 7.205 + 0.213plant-height‡	0.684	0.639	3.788
Above-soil biomass = -27.948 + 5.922soil-pH	0.735	0.697	1.853
Above-soil biomass = 4.484 + 0.118plant-height‡	0.742	0.705	1.827

†For plant height, leaf area and number of leaves, mean values for the five consecutive sampling times (2, 3, 4, 5 and 6 weeks after sowing) were used for the regressions. ‡Plant growth parameters were included among the independent variables. AvP - available phosphorus; Plant-P - plant P content; SEE - standard error of the estimate. All the regressions are significant at $p \leq 0.01$.

lime-manurial amendment option were largely a reflection of the increased availability of soil P and plant uptake of same.

The application of synthetic lime and/or animal manure to acid coarse-textured soils of the humid tropics can influence arable crops production in the region. Increasing the agronomic productivity of these soils would require the use of CaO lime at moderate rates with poultry-droppings manure at about double the optimum 'agronomic' rates. The increases would be due to ameliorated soil acidity and hence elevated soil fertility status evidenced by increased P bioavailability. For the acid sandy-loam soil studied and similar coarse-textured soils of the Derived Savannah, arable crops production would benefit specifically from co-application of CaO lime at 2.5 t ha⁻¹ and poultry-droppings manure at 50 t ha⁻¹. This viable soil fertility management option is, therefore, suggested for coping with soil acidity and related P-fixation problems in arable crops production.

Long-duration field trials on agronomic evaluations of the immediate and residual effects of the suggested option on the overall soil/environmental quality in this and similar agro-environments are, however, needed to comment on its sustainability and validate its adoption. Such future studies can involve various lime and manure types and can explore the relative contributions of organic and inorganic P to the enhanced P bioavailability due to appropriate lime-manure combination.

Acknowledgements

The authors thank the technologists attached to the Teaching & Research Laboratories of the Department of Soil Science at the Faculty of Agriculture of the University of Nigeria, Nsukka for their technical support in the required soil analyses of the study.

References

- Achat D., Bakker M. R., Zeller B., Pellerin S., Benaïme S. and Morel C. 2010. Long-term organic phosphorus mineralization in Spodosols under forests and its relation to carbon and nitrogen mineralization. *Soil Biol. Biochem.*, 42, 1479-1490. <https://doi.org/10.1016/j.soilbio.2010.05.020>
- Adiaha M. S. 2017. The role of organic matter in tropical soil productivity. *World Sci. News*, 86(1), 1-66. <https://www.researchgate.net/publication/351765121>
- Adubasim, C. V., Igwenagu, C. M., Josiah, G. O., Obalum S. E., Okonkwo, U. M., Uzoh, I. M. and Sato, S. 2018. Substitution of manure source and aerator in nursery media on sandy-loam topsoil and their fertility indices 4 months after formulation. *Int. J. Recycl. Org. Waste Agric.*, 7(4), 305-312. <https://dx.doi.org/10.1007/s40093-018-0216-8>
- Agba, O. A., Asiegbu, J. E., Ikenganyia, E. E., Anikwe M. A. N., Omeje, T. E. and Adinde, J. O. 2017. Effects of lime on growth and yield of *Mucuna flagellipes* (Vogel ex Hook) in an acid tropical Ultisol. *J. Agric. Ecol. Res. Int.*, 12(3), 1-14, Article No. JAERI 22741
- Akpan, J. F., Umunnakwe, O. C. and Omedewu, L. I. 2023. Effects of lime and poultry manure on soil properties and cucumber (*Cucumis sativum* L) performance in the humid tropics of southern Nigeria. *Global J. Agric. Sci.*, 22, 101-107. <https://dx.doi.org/10.4314/gjass.v22i1.11>
- Allison, S. D. and Vitousek, P. M. 2005. Responses of extracellular enzymes to simple and complex nutrient inputs. *Soil Biol. Biochem.*, 37, 937-944. <https://doi.org/10.1016/j.soilbio.2004.09.014>
- Ameyu, T. 2019. A review on the potential effect of lime on soil properties and crop productivity improvements. *J. Environ. Earth Sci.*, 9(2), 2225-0948. <https://doi.org/10.7176/JEES>
- Amsalu, S. and Beyene, S. 2020. Effects of lime and phosphorous application on chemical properties of soil, dry matter yield, and phosphorus concentration of barley (*Hordeum vulgare*) grown on Nitisols of Emdibir, Southern Ethiopia. *J. Soil Sci. Environ. Manage.*, 11(4), 131-141. <https://doi.org/10.5897/JSEEM2020.0837>
- Antoniadis, V., Hatzis, F., Bachtsevanidis, D. and Koutroubas, S. D. 2015. Phosphorus availability in low-P and acidic soils as affected by liming and P addition. *Commun. Soil Sci. Plant Anal.*, 46, 1288-1298. <https://doi.org/10.1080/00103624.2015.1033539>
- Bolan, N. S., Adriano, D. C. and Curtin, D. 2003. Soil acidification and liming interactions with nutrient and heavy metal transformation and bioavailability. *Adv.*

- Agron., 78, 215-272. [http://dx.doi.org/10.1016/S0065-2113\(02\)78006-1](http://dx.doi.org/10.1016/S0065-2113(02)78006-1)
- Borggaard, O. K., Raben-Lange, B., Gimsing, A. L. and Strobel B.W. 2005. Influence of humic substances on phosphate adsorption by aluminium and iron oxides. *Geoderma*, 127, 270-279. <https://doi.org/10.1016/j.geoderma.2004.12.011>
- Brady, N. C. and Weil, R. R. 2008. The nature and properties of soils, 14th ed., Pearson, Upper Saddle River, USA
- Bueis, T., Bravo, F., Pando, V., Kissil, Y-A. and Turrión, M-B. 2019. Phosphorus availability in relation to soil properties and forest productivity in *Pinus sylvestris* L. plantations. *Ann For. Sci.*, 76, 97. <https://doi.org/10.1007/s13595-019-0882-3>
- Caires, E. F., Garbuio, F. J., Churka, S., Barth, G. and Corrêa, J. C. L. 2008. Effects of soil acidity amelioration by surface liming on no-till corn, soybean, and wheat root growth and yield. *Eur. J. Agron.*, 28(1), 57-64. <https://doi.org/10.1016/j.eja.2007.05.002>
- Chukwuma, C. C., Oraegbunam, C. J., Ndeshala, S. D., Uchida, Y., Ugwu, V. U., Obalum, S. E. and Igwe, C. A. 2024. Phosphorus mineralization in two lithologically dissimilar tropical soils as influenced by animal manure type and amendment-to-sampling time interval. *Comm. Soil Sci. Plant Anal.*, 55(5), 707-722. <https://doi.org/10.1080/00103624.2023.2276269>
- Dada, B. F. and Ewulo, B. S. 2011. Manure and lime amendment effects on soil pH, fertility, microbe population and respiration. *South Asian J. Exp. Biol.*, 1(4), 172-177; 2011
- Debicka, M, Kocowicz, A., Weber, J. and Jamroz, E. 2016. Organic matter effects on phosphorus sorption in sandy soils. *Arch. Agron. Soil Sci.*, 62(6), 840-855. <https://doi.org/10.1080/03650340.2015.1083981>
- Ebido, N. E., Edeh, I. G., Unagwu, B. O., Nnadi, A.L., Ozongwu, O. V., Obalum S.E. and Igwe, C. A. 2021. Rice-husk biochar effects on organic carbon, aggregate stability and nitrogen-fertility of coarse-textured Ultisols evaluated using *Celosia argentea* growth. *SAINS TANAH – J. Soil Sci. Agroclimatol.*, 18(2), 177-187. <https://dx.doi.org/10.20961/stjsa.v18i2.56330>
- Ejersa, M. T. 2021. Causes of soil acidity and its management mechanisms in Ethiopia: a review. *J. Agric. Sci. Bot.*, 5(4), 054
- Ewulo, B. S., Ojeniyi, S. O. and Akanni, D. A. 2008. Effect of poultry manure on selected soil physical and chemical properties, growth, yield and nutrient status of tomato. *Afr. J. Agric. Res.*, 3(9), 612-616. <https://doi.org/10.5897/AJAR.9000218>
- Fageria, N. K., Baligar, V. C. and Jones, C.A. 2011. Growth and mineral nutrition of field crops. CRC Press, Boca Raton, FL, USA. <https://doi.org/10.1201/b10160>
- Garbuio, F. J., Jones, D. L., Alleoni, L. R., Murphy, D. V. and Caires, E. F. 2011. Carbon and nitrogen dynamics in an Oxisol as affected by liming and crop residues under no-till. *Soil Sci. Soc. Am. J.*, 75, 1723-1730. <https://doi.org/10.2136/sssaj2010.0291>
- Gichangi, E. 2019. Effects of organic amendments on the transformations and bioavailability of phosphorus in soils: A review. *Discov. Agric.*, 5, 41-50. https://www.discoveryjournals.org/agriculture/current_issue/2019/A5.pdf
- Grover, S. P., Butterly, C. R., Wang, X. and Tang, C. 2017. The short-term effects of liming on organic carbon mineralization in two acidic soils as affected by different rates and application depths of lime. *Biol. Fert. Soils*, 53, 431-443. <https://doi.org/10.1007/s00374-017-1196-y>
- Hanč, A., Tlustoš, P., Száková, J. and Balík, J. 2008. The influence of organic fertilizers application on phosphorus and potassium bioavailability. *Plant, Soil Environ.*, 54(6), 247-254
- Haynes, R. J. and Mokolobate, M. S. 2001. Amelioration of Al toxicity and P deficiency in acid soils by additions of organic residues: a critical review of the phenomenon and the mechanisms involved. *Nutr. Cycl. Agroecosystems*, 59, 47-63. <https://doi.org/10.1023/A:1009823600950>
- Ifeanyi-Onyishi, I. F., Ezeaku, V. I., Umeugokwe, C. P., Ezeaku, P. I. and Obalum, S. E. 2024. Distribution of soil fertility indices in aggregate size fractions under different land-use types for coarse-textured soils of the derived savannah. *J. Degraded Mining Lands Manage.*, 12(1), 6695-6704. <http://doi.org/10.15243/jdmlm.2024.121.6695>
- Islam, M. R., Jahan, R., Uddin, S., Harine, I.J., Hoque M. A., Hassan, S., Hassan, M. M. and Hossain, M. A. 2021. Lime and organic manure amendment enhances crop productivity of wheat-mungbean-T. aman cropping pattern in acidic Piedmont soils. *Agronomy*, 11, 1595. <https://doi.org/10.3390/agronomy11081595>
- Kabango, F., John, B., Onesimus, S., Lenssen, A. W., Majaliwa M.J.G. and Stewart K. 2022. Effect of applying low rates of lime and manure on bean growth and yield on Ferralsols of Lake Victoria crescent agro-ecological Zone-Central Uganda. *Afr. J. Agr. Res.*, 18(12), 1110-1120. <https://doi.org/10.5897/AJAR2022.16176>
- Li Y., Cui S., Chang, S. X. and Zhang, Q. 2019. Liming effects on soil pH and crop yield depend on lime

- material type, application method and rate, and crop species: a global meta-analysis. *J. Soils Sediments*, 19, 1393-1406. <https://doi.org/10.1007/s11368-018-2120-2>
- Materechera, S. A. and Mkhabela, T. S. 2002. The effectiveness of lime, chicken manure and leaf litter ash in ameliorating acidity in a soil previously under black wattle (*Acacia mearnsii*) plantation. *Bioresour. Technol.* 85, 9-16
- McLean, E. O. 1982. Soil pH and lime requirement. In: Page, A. L., Miller, R. H. and Keeney D. (eds.), *Methods of soil analysis, Part 2: chemical methods*, 2nd edn. Agronomy Monograph No. 9 (pp. 199-224), Am. Soc. Agron., Madison Wisconsin
- Mkhonza, N. P., Buthelezi-Dube N.N. and Muchaonyerwa P. 2020. Effects of lime application on nitrogen and phosphorus availability in humic soils. *Scientific Reports*, 10: 8634. <https://doi.org/10.1038/s41598-020-65501-3>
- Moreira, A., Sfredo, G. J., Moraes, L. A. C. and Fageria N. K. 2015. Lime and cattle manure in soil fertility and soybean grain yield cultivated in tropical soil. *Commun. Soil Sci. Plant Anal.*, 46(9), 1157-1169. <https://doi.org/10.1080/100103624.2015.1033542>
- Ndzeshala, S. D., Obalum, S. E. and Igwe, C. A. 2021. Effect of lime and phosphorous sources on soil fertility, nutrient uptake, growth and yield of soybean (*Glycine max*) in Southeastern Nigeria. In: *Understanding Soil Organic Matter Dynamics: Key to Sustainable Ecosystem Health (Book of Abstracts, Abstract C044)*, the 45th Annual Conference of Soil Science Society of Nigeria (SSSN), 17-21 May 2021, Bowen University, Iwo, Osun State, Nigeria
- Ndzeshala, S. D., Obalum, S. E. and Igwe, C. A. 2023. Some utilisation options for cattle dung as soil amendment and their effects in coarse-textured Ultisols and maize growth. *Int. J. Recycl. Org. Waste Agric.*, 12(1), 123-39. <https://doi.org/10.30486/ijrowa.2022.1934239.1284>
- Nnadi, A. L., Nnanna, P. I., Onyia, V. N., Obalum, S. E. and Igwe, C. A. 2020. Growth and yield responses of high-density coverage sweet potato to liming and fertilizer combinations for sandy-loam Ultisols at Nsukka, southeastern Nigeria. In: *Climate- Smart Soil Management, Soil Health/ quality and Land Management Synergies for Sustainable Ecosystem Services* (pp 263-269), Proceedings of the 44th Conference of Soil Science Society of Nigeria. Colloquia SSSN 44, 263-269. <https://doi.org/10.36265/colssn.2020.4438>
- Nnadi, A. L., Ugwu, V. U., Nwite, J. C., Obalum, S. E., Igwe, C. A. and Wakatsuki, T. 2021. Manurial amendments and source of water for supplemental irrigation of sawah-rice system influenced soil quality and rice yield. *Agro-Science*, 20(1), 95-102. <https://dx.doi.org/10.4314/as.v20i1.15>
- Nwite, J. C., Keke, C. I., Obalum, S. E., Essien, J. B., Anaele, M. U. and Igwe, C. A. 2013. Organo-mineral amendment options for enhancing soil fertility and nutrient composition and yield of fluted pumpkin. *Int. J. Veg. Sci.*, 19(2), 188-199. <https://doi.org/10.1080/19315260.2012.705233>
- Obalum, S. E. and Chibuike, G. U. 2017. Air-drying effect on soil reaction and phosphorus extractability from upland-lowland tropical soils as related to their colloidal stability. *Appl. Ecol. Environ. Res.*, 15(1), 525-540. OnlineFirst accessible at: <http://aloki.hu/indvolOF.htm>
- Obalum, S. E. and Obi, M. E. 2014. Measured versus estimated total porosity along structure-stability gradients of coarse-textured tropical soils with low-activity clay. *Environ Earth Sci.*, 72, 1953-1963. <https://doi.org/10.1007/s12665-014-3102-3>
- Obalum, S. E., Igwe, C. A., Obi, M. E. and Wakatsuki, T. 2011b. Water use and grain yield response of rainfed soybean to tillage-mulch practices in southeastern Nigeria. *Scientia Agricola*, 68(5), 554-561. <https://doi.org/10.1590/S0103-90162011000500007>
- Obalum, S. E., Nwite, J. C., Watanabe, Y., Igwe, C. A. and Wakatsuki, T. 2012. Comparative topsoil characterization of sawah rice fields in selected inland valleys around Bida, north-central Nigeria: physicochemical properties and fertility status. *Trop. Agric.Dev.*, 56(2), 39-48. <https://doi.org/10.11248/jsta.56.39>
- Obalum, S. E., Ofem, K. I., Nwamba, D. C., Joseph, P. O., Edeh, I. G., Amalu, U. C. and Igwe, C. A. 2024. Aggregate stability responses of three derived-savannah soils to poultry-droppings manure at different pulverisation-to-sampling time intervals. *Appl. Environ. Biotechnol.*, 9(1), 38-51. <http://doi.org/10.26789/AEB.2024.01.005>
- Obalum, S. E., Ogumba, P. O. and Uzoh, I. M. 2020. Influence of tillage-seedbed and manure-NPK-micronutrient management options on selected soil properties of sandy-loam Ultisols evaluated using sweet potato. *Nigerian J. Soil Sci.*, 30(3), 117-125
- Obalum, S. E., Okpara, I. M., Obi, M. E. and Wakatsuki, T. 2011a. Short-term effects of tillage-mulch practices under sorghum and soybean on organic carbon and eutrophic status of a degraded Ultisol in southeastern Nigeria. *Trop. Subtrop. Agroecosyst.*, 14(2), 393-403
- Ogumba P.O., Orah A.I., Ndzeshala S.D., Aneke A.M., Nnadi A.L., Law-Ogbomo K.E., Obalum S.E. and Igwe

- C. A. 2024. Synthetic lime and manure-NPK effects in sandy-loam Ultisols after growing sweet potato in successive rainy and dry seasons. *Agriculturae Conspectus Scientificus* (submitted)
- Ogunzei, K. C., Okebalama, C. B. and Obalum, S. E. 2019. Optimum poultry droppings rate for coarse-loamy Ultisols based on soil macro-aggregation and fertility indices and evaluation using cucumber (*Cucumis sativus*). Presented under Understanding Nigerian Soils for Sustainable Food & Nutrition Security and Healthy Environment, the 43rd Conference of Soil Science Society of Nigeria on (SSSN), 15-19 July 2019, University of Agriculture, Makurdi, Benue State, Nigeria
- Okebalama, C. B., Ibezim, C. O., Ugwu, V. U. and Marschner, B. 2020. Effects of amending an Ultisol with bambara groundnut seed residue on soil fertility, microbial nutrient cycling and yield of cucumber. *Biol. Agric. Hortic.*, 36(4), 249-266. <https://doi.org/10.1080/01448765.2020.1808067>
- Olsen, S. R. and Sommers, L. E. 1982. Phosphorous. In: Page, A. L., Miller, R. H. and Keeney, D. R. (eds.), *Methods of soil analysis, Part 2: chemical properties*, 2nd edn. Agronomy Monograph No. 9 (pp.15-72), Am. Soc. Agron., Madison Wisconsin
- Onah, C. J., Nnadi, A. L., Eyibio, N. U., Obi, J. O., Orah, A. I., Amuji, C. F. and Obalum, S. E. 2023. Off-season heavy application of poultry manure to droughty-acid soils under heavily protective organic mulch later burnt to ash improves their productivity. *West Afr. J. Appl. Ecol.*, 31(1), 23-36
- Osakwe, U. C., Nnadi, A. L., Ede, L. C., Obi, J. O., Okolo, C. C. and Obalum, S. E. 2023. Surface mulching effects on soil temperature of jumbo-size potted coarse-textured Ultisols and evaluation on sorghum and soybean growth. *Agro-Science*, 22(4), 82-90. <https://dx.doi.org/10.4314/as.v22i4.12>
- Opala, P. A. 2017. Influence of lime and phosphorus application rates on growth of maize in an acid soil. *Adv. Agric.*, 3, 1-5. <https://doi.org/10.1155/2017/7083206>
- Paradelo, R., Virto, I. and Chenu, C. 2015. Net effect of liming on soil organic carbon stocks: a review. *Agric. Ecosyst. Environ.* 202, 98-107. <https://doi.org/10.1016/j.agee.2015.01.005>
- Rastija, D., Zebec, V. and Rastija, M. 2014. Impacts of liming with dolomite on soil pH and phosphorus and potassium availabilities. *Novenytermeles*, 63, 193-196. <https://api.semanticscholar.org/CorpusID:96714913>
- Rousk, J., Brookes, P. C. and Baath, E. 2010. Investigating the mechanisms for the opposing pH relationships of fungal and bacterial growth in soil. *Soil Biol. Biochem.*, 42, 926-934. <https://doi.org/10.1016/j.soilbio.2010.02.009>
- Saleem, A., Irshad, M., Hassan, A., Mahmood, Q. and Eneji A. E. 2017. Extractability and bioavailability of phosphorus in soils amended with poultry manure co-composted with crop wastes. *J. Soil Sci. Plant Nutr.*, 17(3), 609-623. <http://dx.doi.org/10.4067/S0718-95162017000300005>
- Sandrakirana, R. and Arifin, Z. 2021. Effect of organic and chemical fertilizers on the growth and production of soybean (*Glycine max*) in dry land. *Rev. Fac. Nac. Agron. Medellín*, 74(3), 9643-9653. <https://doi.org/10.15446/rfnam.v74n3.90967>
- Sato, S. and Comerford, N. B. 2005. Influence of soil pH on inorganic phosphorus sorption and desorption in a humid Brazilian Ultisol. - *Revista Brasileira de Ciência do Solo*, 29, 685-694. <https://doi.org/10.1590/S0100-06832005000500004>
- Soremi, A. O., Adetunji, M. T., Adejuyigbe, C. O., Bodunde J.G. and Azeez J.O. 2017. Effects of poultry manure on some soil chemical properties and nutrient bioavailability to soybean. *J. Agric. Ecol. Res. Int.*, 11(3), 1-10. <http://dx.doi.org/10.9734/JAERI/2017/32419>
- Stammer, A. J. and Mallarino, A. P. 2018. Plant tissue analysis to assess phosphorus and potassium nutritional status of corn and soybean. *Soil Sci. Soc. Am. J.*, 82, 260-270. <https://doi.org/10.2136/sssaj2017.06.0179>
- Takala, B. 2019. Soil acidity and its management options in western Ethiopia: review. *J. Environ. Earth Sci.*, 9(10), 225-0948
- Desalegn, T., Alemu G., Adella A., Debele T. and Gonzalo-J J. 2016. Effect of lime and phosphorus fertilizer on acid soils and barley (*Hordeum vulgare* L.) performance in the central highlands of Ethiopia. *Exp. Agric.*, 53, 432-444. <http://dx.doi.org/10.1017/S0014479716000491>
- Tshiabukole, J. P. K., Khonde, G. P., Phongo, A. M., Ngoma, N., Vumilia, R. K. and Kankolongo, A. M. 2022. Liming and mineral fertilization of acid soils in maize crop within the Savannah of Southwestern of Democratic Republic of Congo. *Open Access Libr. J.*, 9, 8412. <https://doi.org/10.4236/oalib.1108412>
- Turrión, M-B., Schneider, K. and Gallardo, J. F. 2008. Soil P availability along a catena located at the Sierra de Gata Mountains, Western Central Spain. *For. Ecol. Manag.*, 255, 3254-3262 <https://doi.org/10.1016/j.foreco.2008.01.076>
- Ubi, M. W., Ubi, W. and Ubi, G. M. 2017. Effect of lime application on soil physicochemical characteristics and plant nutrient contents of three soybean cultivars

- [*Glycine max* (L.) Merrill] in coastal plain sands of Nigeria. Direct Res. J. Agric. Food Sci., 5(2), 96-106. <https://directresearchpublisher.org/drjafs/files/2017/01/Ubi-et-al.pdf>
- Ugwu, D. O., Joshua, P. E., Obalum, S. E., Dedan, N. K. and Njoku, O. U. 2024. Aging-associated fermentation of palm oil-mill effluent enhances its organo-fertilizer value and the desired agronomic effects in low-fertility soils. Int. J. Recycl. Org. Waste Agric., 13(4), 132439. <https://dx.doi.org/10.57647/ijrowa-xqpd-6789>
- Ugwu, V. U., Nnadi, A. L., Adubasim, C. V., Sato, S., Igwenagu, C. M., Obalum, S. E. and Igwe, C. A. 2020. Organic-waste aerator could completely displace poultry-droppings manure in nursery media based on coarse-textured soil: evidence with cashew seedlings. In: Baiyeri KP and Aba SC (eds), Sustainable Horticulture Production System Intensified (pp. 941-951), Proceedings of the 38th Annual Conference of the Horticultural Society of Nigeria (HORTSON), University of Nigeria Nsukka, 25-31 October 2020
- Ukabalala, M. E., Kolo, J., Obalum, S. E., Amhakhian, S. O., Igwe C.A. and Hermensah. 2021. Physicochemical properties as related to mineralogical composition of floodplain soils in humid tropical environment and the pedological significance. Environ. Monit. Assess. 193:569. <https://doi.org/10.1007/s10661-021-09329-y>
- Umeugokwe, C. P., Ugwu, V. U., Umeugochukwu, O. P., Uzoh, I. M., Obalum, S. E., Ddamulira, G., Karwani, G. M. and Alenoma, G. 2021. Soil fertility indices of tropical loamy sand as influenced by bambara groundnut variety, plant spacing and fertilizer type. Agro-Science, 20(1), 65-71. <https://dx.doi.org/10.4314/as.v20i1.11>
- Verde, B. S., Danga, B. O. and Mugwe, J. N. 2013. Effects of manure, lime and mineral P fertilizer on soybean yields and soil fertility in a humic nitisol in the Central Highlands of Kenya. Int. J. Agric. Sci. Res., 2(9), 283-291
- Wang, X. R., Shen, J. B. and Liao, H. 2010. Acquisition or utilization, which is more critical for enhancing phosphorus efficiency in modern crops? Plant Sci., 179, 302-306. <https://doi.org/10.1016/j.plantsci.2010.06.007>
- Zhang, P., Zhong, K., Tong, H., Shahid, M. Q. and Li, J. 2016. Association mapping for aluminum tolerance in a core collection of rice landraces. Fron. Plant Sci., 7, 1415. <https://doi.org/10.3389/fpls.2016.01415>
-