

Evaluation of Managing Nutrient Supply to Soybean Crop on Calcareous Soil

G. R. Nirmal, S. M. Todmal, B. M. Kamble, M. R. Chauhan, A. G. Durgude

Department of Soil Science, Mahatma Phule Krishi Vidyapeeth, Rahuri – 413 722 (India)

Corresponding author Email : gaurinirmal789@gmail.com

(Received : 20.08.2025 Accepted : 21.11.2025)

Abstract

Calcareous soils, common in arid and semi-arid regions, are characterised by high calcium carbonate content, alkaline pH, and limited availability of macro and micronutrients. These constraints often reduce crop performance, particularly in soybean (*Glycine max* L.), which is sensitive to nutrient imbalances and iron chlorosis. This study investigated the effect of integrated nutrient management strategies combining inorganic fertilizers, sulphur application, organic amendments, and foliar nutrient sprays on soil properties in calcareous soils. It was aimed to study effect on nutrient management on soybean on calcareous soil using a Completely Randomized Design (CRD) comprised of seven treatments *viz.*; T₁ - Absolute control, T₂ - RDF (50:75:45 N: P₂O₅: K₂O kg ha⁻¹), T₃ - RDF + two foliar spray of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS, T₄ - RDF (50 kg N through SCU + 75:45 P₂O₅: K₂O kg ha⁻¹) + two foliar spray of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS, T₅ - RDF + 40 kg S ha⁻¹ + two foliar spray of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS, T₆ - RDF + soil application of micro-grade I at sowing @ 25 kg ha⁻¹ + two foliar spray of 19:19:19 (1%) at 30 and 55 DAS and T₇ - RDF + 40 kg S ha⁻¹ + soil application of micro-grade I at sowing @ 25 kg ha⁻¹ + two foliar spray of 19:19:19 (1%) at 30 and 55 DAS replicated 3 times. Post-harvest soil analysis revealed improved soil available nutrients *viz.*, available N (148 kg ha⁻¹), available P (17.33 kg ha⁻¹), available K (286 kg ha⁻¹) and available S (12.21 mg kg⁻¹), higher organic carbon (0.45%), and a reduction in calcium carbonate content (11.21%). The findings highlight that balanced nutrient supply, incorporating both macro- and micronutrients through combined application methods, can effectively overcome the limitations of calcareous soils, improve soybean productivity and sustain soil health.

Key words : Nutrient management, calcareous soil, soybean.

Calcareous soils, covering about 30% of global land and widely found in Indian states such as Maharashtra, Rajasthan, and Uttar Pradesh, contain >15% calcium carbonate with pH >7.5, which limits nutrient solubility and availability. Such conditions cause micronutrient deficiencies, soil crusting, and reduced crop productivity. Soybean (*Glycine max* L.), an important oilseed and protein-rich legume, is highly sensitive to these constraints, often showing Fe and Zn deficiencies that result in chlorosis, poor growth, and yield loss. INM combining fertilizers, organics, secondary nutrients, and foliar sprays, offers a sustainable strategy to improve nutrient availability, crop performance, and soil fertility.

Study area : Geographically the central campus of MPKV, Rahuri is situated between 19°34' N latitude and 74°64' E longitude elevation of 513 meter above Mean Sea Level.

Material and Methods

The pot culture experiment was conducted in the Wire House of the Department of Soil Science, Post Graduate Institute, MPKV, Rahuri. Seven treatments *viz.*; T₁ - Absolute control, T₂ - RDF (50:75:45 N: P₂O₅: K₂O kg ha⁻¹), T₃ - RDF + two foliar spray of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS, T₄ - RDF (50 kg N through SCU + 75:45 P₂O₅: K₂O kg ha⁻¹) + two foliar spray of [19:19:19 (1%) + Phule liquid micro-grade II

(0.5%)] at 30 and 55 DAS, T_5 - RDF + 40 kg S ha^{-1} + two foliar spray of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS, T_6 - RDF + soil application of micro-grade I at sowing @ 25 kg ha^{-1} + two foliar spray of 19:19:19 (1%) at 30 and 55 DAS and T_7 - RDF + 40 kg S ha^{-1} + soil application of micro-grade I at sowing @ 25 kg ha^{-1} + two foliar spray of 19:19:19 (1%) at 30 and 55 DAS were given to soybean.

Plastic pots with a capacity of 20 kg were used. The calcareous soil for the experiment was collected from the Micronutrient Research Farm. Each pot was filled with 20 kg of soil, leaving two to three inches portion at the top to facilitate watering.

Soybean seeds of variety Phule Sangam (KDS 726) were used. The sowing was done with dibbling 5 seeds per pot at appropriate spacing and then thinning was done at three-leaf stage of crop, leaving two healthy seedlings per pot. The experiment was laid in Completely Randomized Design (CRD) with seven treatments each replicated three times.

The hydrogen ion activity (pH) of soil samples was determined by using a pH meter and electrical conductivity by conductivity bridge by using 1:2.5 soil: water suspension (Jackson, 1973). Soil organic carbon content was determined by Walkley and Black's wet oxidation method as described by Piper (1966). Calcium carbonate content of soil was determined by using rapid titration method as suggested by Puri (1949). Soil available nitrogen content in the soil was determined by alkaline potassium permanganate method (Subbiah and Asija, 1956). Soil available phosphorus was determined by Olsen's method using 0.5 M sodium bicarbonate as an extractant. Intensity of blue colour developed was measured by using spectrophotometer at 882 nm wavelength (Watanabe and Olsen, 1965). Soil available

potassium was determined by using neutral normal ammonium acetate as an extractant using flame photometer (Jackson, 1973). Soil available sulphur was determined on spectrophotometer by using calcium chloride as an extractant (Williams and Steinberg, 1959)

Result and discussion

The experiment was conducted at central campus MPKV, Rahuri. The result of the investigation is described under following heading

Initial status of soil : The soil was alkaline in reaction having pH (8.40) with low salt content (0.29 dS m^{-1}). Soil organic carbon content was low (0.35%) and calcium carbonate content was high (12.23%). The soil was low in available N (136 kg ha^{-1}), low in available P (13.5 kg ha^{-1}), high in available K (217 kg ha^{-1}) and medium in available S (10.36 kg ha^{-1}). Whereas, the soil was sufficient in manganese and was deficient in iron, zinc and copper

Effect of nutrient management on soil chemical properties : Details of data regarding chemical properties of calcareous soil after harvest of soybean is presented in Table 1.

Soil pH and Electrical conductivity : The effect of nutrient management on soil chemical properties *viz.*, soil pH and electrical conductivity were found non-significant.

Soil organic carbon : The treatment T_7 recorded significantly highest soil organic carbon (0.45%) and it was followed by treatment T_5 (0.44%).

Calcium carbonate content : The calcium carbonate content showed a slight decreasing trend with nutrient management. The lowest calcium carbonate (11.21%) was recorded in treatment T_7 .

Effect of nutrient management on soil available macronutrients : Details of data regarding available nutrients in calcareous soil after harvest of soybean are presented in

Table 2.

Soil available nitrogen : The soil available nitrogen (148 kg ha⁻¹) content at harvesting of

Table 1. Effect of nutrient management on chemical properties of soil after harvest of soybean

Treatment	pH (dS m ⁻¹)	EC (%)	OC (%)	CaCO ₃
T ₁ - Absolute control	8.40	0.30	0.36	12.17
T ₂ - RDF (50:75:45 N: P ₂ O ₅ : K ₂ O kg ha ⁻¹)	8.38	0.34	0.39	12.10
T ₃ - RDF + Two foliar sprays of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS	8.37	0.33	0.42	11.85
T ₄ - RDF (50 kg N through SCU + 75:45 P ₂ O ₅ : K ₂ O kg ha ⁻¹) + Two foliar sprays of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS	8.36	0.33	0.43	11.79
T ₅ - RDF + 40 kg S ha ⁻¹ + Two foliar sprays of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS	8.32	0.36	0.44	11.46
T ₆ - RDF + Soil application of micro-grade I at sowing @ 25 kg ha ⁻¹ + Two foliar sprays of 19:19:19 (1%) at 30 and 55 DAS	8.36	0.35	0.41	11.67
T ₇ - RDF + 40 kg S ha ⁻¹ + Soil application of micro-grade I at sowing @ 25 kg ha ⁻¹ + Two foliar sprays of 19:19:19 (1%) at 30 and 55 DAS	8.35	0.35	0.45	11.21
Mean	8.36	0.34	0.41	11.75
Initial	8.40	0.29	0.35	12.23
SE (m) ±	0.01	0.01	0.02	0.18
C.D. (5%)	NS	NS	0.05	0.57

Table 2. Effect of nutrient management on available nutrients in soil after harvest of soil

Treatment	Available nutrients			
	N	P	K	S
	(kg ha ⁻¹)			(mg kg ⁻¹)
T ₁ - Absolute control	132	11.90	229	10.24
T ₂ - RDF (50:75:45 N: P ₂ O ₅ : K ₂ O kg ha ⁻¹)	138	13.18	261	10.53
T ₃ - RDF + Two foliar sprays of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS	140	14.65	257	10.58
T ₄ - RDF (50 kg N through SCU + 75:45 P ₂ O ₅ : K ₂ O kg ha ⁻¹) + Two foliar sprays of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS	148	15.50	238	11.24
T ₅ - RDF + 40 kg S ha ⁻¹ + Two foliar sprays of [19:19:19 (1%) + Phule liquid micro-grade II (0.5%)] at 30 and 55 DAS	141	16.70	286	12.12
T ₆ - RDF + Soil application of micro-grade I at sowing @ 25 kg ha ⁻¹ + Two foliar sprays of 19:19:19 (1%) at 30 and 55 DAS	144	14.78	241	11.32
T ₇ - RDF + 40 kg S ha ⁻¹ + Soil application of micro-grade I at sowing @ 25 kg ha ⁻¹ + Two foliar sprays of 19:19:19 (1%) at 30 and 55 DAS	146	17.33	240	12.21
Mean	142	14.86	250	11.18
Initial	136	13.50	217	10.36
SE (m) ±	2.72	0.76	14.70	0.43
C.D. (5%)	8.25	2.33	44.60	1.30

soybean was significantly highest in treatment T₄. There was improvement in soil available nitrogen in all nutrient management treatments.

Soil available phosphorus : The soil available phosphorus (17.33 kg ha⁻¹) was found significantly highest in treatment T₇ and it was at par with the treatment T₅ (16.70 kg ha⁻¹) and the treatment T₄ (15.50 kg ha⁻¹).

Soil available potassium : The significantly maximum soil available potassium (286 kg ha⁻¹) was recorded in treatment T₅, it was followed by treatment T₂ (261 kg ha⁻¹) and the treatment T₃ (257 kg ha⁻¹).

Soil available sulphur : The soil available sulphur (12.21 mg kg⁻¹) was found significantly highest in treatment T₇ and it was at par with the treatment T₅ (12.12 mg kg⁻¹), the treatment T₆ (11.32 mg kg⁻¹) and the treatment T₄ (1.24 mg kg⁻¹).

Conclusion

Integrated nutrient management involving RDF, soil-applied sulphur, and foliar sprays of balanced macro- and micronutrients substantially improved soil fertility, reduced calcium carbonate levels, and promoted sustainable nutrient cycling. Adoption of such integrated practices can significantly boost soybean production in calcareous regions while preserving soil health.

Reference

- Bolan, N., Srivastava, P., Rao, C. S., Satyanaraya, P. V., Anderson, G. C., Bolan, S., ... and Kirkham, M. B. 2023. Distribution, characteristics and management of calcareous soils. *Advances in agronomy*, 182, 81-130.
- Jackson, M. L. 1973. *Soil chemical analysis*. Prentice Hall of India, New Delhi. pp. 498.
- Piper, C. S. 196.) *Soil and Plant Analysis*, Hans, Publication Bombay, Asian (Edition) pp-368.
- Puri, A. H. 1949. *Soils, Their Physics and Chemistry*, Reinhold Publ. Corporation New York U.S.A.
- Subbiah, B. V., and Asija, A. L. 1956. A rapid procedure for the estimation of available Nitrogen in soil. *Current Science* 25, 259-260.
- Taalab, A. S., Ageeb, G. W., Siam, H. S., and Mahmoud, S. A. 2019. Some characteristics of calcareous soils. *A Review* 8(1), 96-105.
- Thapa, S., Bhandari, A., Ghimire, R., Xue, Q.z Kidwaro, F., Ghatrehsamani, S., ... and Goodwin, M. 2021. Managing micronutrients for improving soil fertility, health and soybean yield. *Sustainability*, 13(21), 11766.
- Wahba, M., Fawkia, L. A. B. I. B., and Zaghloul, A. 2019. Management of calcareous soils in arid region. *International Journal of Environmental Pollution and Environmental Modelling*, 2(5), 248-258.
- Watanabe, F. S. and Olsen, S. R. 1965. Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from the soil. *Soil Science Society of America Journal*, 29, 677-678.
- Williams, C. H. and Steinberg, A. 1959. Sulphur fraction as chemical indices of available sulphur in some Australian Soil. *Australian Journal of Agriculture Research*. 10, 340-350.